

ATTACHMENT A (ICD-10)
ANTIPSYCHOTIC AGENTS
For KEPRO Use Only

Appropriate Diagnoses for All Antipsychotics:

Diagnoses	ICD-10 Codes
Autistic disorder	F84.0
Bipolar I disorder	F31.0,F31.1,F31.6
Bipolar II disorder	F31.81
Bipolar disorder not otherwise specified	F31.9
Conduct disorder	F91.1,F91.2,F91.9
Delusional disorder	F22
Dysthymic disorder	F34.1,F41.8
Major depressive disorder	F32,F33
Mood disorder not otherwise specified	F39
Schizophrenic disorders	F20
Tourette's disorder	F95.2
Unspecified mental disorders and psychosis	F09,F29

Additional Drug Specific Diagnoses for All Ages:

Diagnoses	ICD-10 Codes
Disorders of porphyrin metabolism (chlorpromazine)	E80.0,E80.1,E80.2
Intractable hiccups (chlorpromazine)	F45.8, R06.6
Nausea and vomiting (chlorpromazine and perphenazine)	G43.A0,R11
Nonpsychotic anxiety (trifluoperazine)	Z63.7,Z63.71,Z63.72,Z63.79

Additional Diagnoses and Targeted Behaviors¹ for Children:

Diagnoses	ICD-10 Codes
Explosive aggression in conduct disorder	F63.81,F91.1,F91.2
Disruptive behaviors and agitation in youngsters with developmental delay (mental retardation, autism, or pervasive developmental disorders)	F84,F89,2,Z72.810,F91.9,F70-F79
Tic disorders and Tourette's syndrome	F95.2,F95.9
Adolescent-onset bipolar disorders	F31.0,F31.1,F31.6, F31.81,F31.9
Childhood and adolescent onset schizophrenia	F20
Early onset psychotic depression	F32,F33
Disruptive behaviors and agitation in youngsters with traumatic brain injury	F91.9,S0.60,S06.2,S06.3,S06.4,S06.9

¹Source: Martin A, Scahill L, Charney D, Leckman J. Pediatric Psychopharmacology Principles and Practice. Oxford University Press 2003;149.